


# The Pumping Lemma for Context-Free Languages

Take an **infinite** context-free language



Generates an infinite number  
of different strings

Example:

$$S \rightarrow AB$$

$$A \rightarrow aBb$$

$$B \rightarrow Sb$$

$$B \rightarrow b$$

$$S \rightarrow AB$$

$$A \rightarrow aBb$$

$$B \rightarrow Sb$$

$$B \rightarrow b$$

In a derivation of a long string,  
variables are repeated

A derivation:


$$S \Rightarrow AB \Rightarrow aBbB \Rightarrow abb\underline{B}$$

$$\Rightarrow abbSb \Rightarrow abbABb \Rightarrow abbaBbBb \Rightarrow$$

$$\Rightarrow abbabb\underline{B}b \Rightarrow abbabbbb$$


Derivation tree

string *abbabbbb*




Derivation tree

string *abbabbbb*



repeated

$B \Rightarrow Sb \Rightarrow ABb \Rightarrow$   
 $\Rightarrow aBbBb \Rightarrow aBbbb$




\*

$B \Rightarrow aBbbb$

$B \Rightarrow b$

# Repeated Part




\*

$$B \Rightarrow aBbbb$$

Another possible derivation from  $B$

\*  
 $B \Rightarrow aBbbb$




\* \*  
 $B \Rightarrow aBbbb \Rightarrow aaBbbbbbb$







\*  
 $S \Rightarrow abbBb$   
 \*  
 $B \Rightarrow aBbbb$   
 $B \Rightarrow b$



\*  
 $S \Rightarrow abb(a)^2 B(bbb)^2 \Rightarrow abb(a)^2 b(bbb)^2$   
 \*

$$\begin{array}{ccc} * & * & \\ S \Rightarrow abbBb & B \Rightarrow aBbbb & B \Rightarrow b \end{array}$$

---



$$* \\ S \Rightarrow abb(a)^2 b(bbb)^2$$




$$abb(a)^2 b(bbb)^2 \in L(G)$$

$$* \\ S \Rightarrow abbBb$$

$$* \\ B \Rightarrow aBbbb$$

$$B \Rightarrow b$$




$$* \\ S \Rightarrow abb(a)^2 B(bbb)^2 \Rightarrow abb(a)^3 B(bbb)^3$$

$$* \\ S \Rightarrow abbBb$$

$$* \\ B \Rightarrow aBbbb$$

$$B \Rightarrow b$$



$$* \\ S \Rightarrow abb(a)^3 B(bbb)^3 \Rightarrow abb(a)^3 b(bbb)^3$$

$$\begin{array}{ccc} * & * & \\ S \Rightarrow abbBb & B \Rightarrow aBbbb & B \Rightarrow b \end{array}$$

---




$$* \\ S \Rightarrow abb(a)^3 b(bbb)^3$$




$$abb(a)^3 b(bbb)^3 \in L(G)$$

## In General:

$$\begin{array}{ccc} * & * & \\ S \Rightarrow abbBb & B \Rightarrow aBbbb & B \Rightarrow b \end{array}$$


$$* \\ S \Rightarrow abb(a)^i b(bbb)^i$$


$$abb(a)^i b(bbb)^i \in L(G) \quad i \geq 0$$



Consider now an infinite  
context-free language  $L$

Let  $G$  be the grammar of  $L - \{\lambda\}$

Take  $G$  so that It has no unit-productions  
no  $\lambda$ -productions

Let  $p =$  (Number of productions)  $\times$   
(Largest right side of a production,  
let it be  $f$ )

Let  $m = p + 1$

Example  $G: S \rightarrow AB$

$A \rightarrow aBb$

$B \rightarrow Sb$

$B \rightarrow b$

$$p = 4 \times 3 = 12$$

$$m = p + 1 = 13$$

Take a string  $w \in L(G)$   
with length  $|w| \geq m$

We will show:

in the derivation of  $w$   
a variable of  $G$  is repeated

$$S \stackrel{*}{\Rightarrow} w$$

$$v_1 \Rightarrow v_2 \Rightarrow \cdots \Rightarrow v_k \Rightarrow w$$

$$S = v_1$$

$$v_1 \Rightarrow v_2 \Rightarrow \cdots \Rightarrow v_k \Rightarrow w$$

$$|v_{i+1}| < |v_i| + f \quad \leftarrow \text{maximum right hand side of any production}$$



$$|w| < k \cdot f$$




$$m \leq |w| \leq k \cdot f \quad \longrightarrow \quad p < k \cdot f$$

$$v_1 \Rightarrow v_2 \Rightarrow \cdots \Rightarrow v_k \Rightarrow w$$

$$p < k \cdot f$$




$$k > \frac{p}{f}$$



Number of productions  
in grammar

$$v_1 \Rightarrow v_2 \Rightarrow \cdots \Rightarrow v_k \Rightarrow w$$

$k >$  Number of productions  
in grammar



Some production must be repeated

$$v_1 \Rightarrow \cdots \Rightarrow a_1 A a_2 \Rightarrow \cdots \Rightarrow a_3 A a_4 \Rightarrow \cdots \Rightarrow w$$

Repeated  
variable

$$\begin{array}{l} S \rightarrow r_1 \\ A \rightarrow r_2 \\ B \rightarrow r_2 \\ \dots \end{array}$$

$$w \in L(G) \quad |w| \geq m$$


Derivation of string  $w$

$$S \Rightarrow \cdots \Rightarrow a_1 A a_2 \Rightarrow \cdots \Rightarrow a_3 A a_4 \Rightarrow \cdots \Rightarrow w$$

Some variable is repeated



# Derivation tree of string $w$



$w = uvxyz$

$u, v, x, y, z :$


Strings of terminals

Possible derivations:

$$S \stackrel{*}{\Rightarrow} uAz$$

$$A \stackrel{*}{\Rightarrow} vAy$$

$$A \stackrel{*}{\Rightarrow} x$$



We know:

$$S \overset{*}{\Rightarrow} uAz$$

$$A \overset{*}{\Rightarrow} vAy$$

$$A \overset{*}{\Rightarrow} x$$

This string is also generated:

$$S \overset{*}{\Rightarrow} uAz \overset{*}{\Rightarrow} uxz$$

$$uv^0xy^0z$$

We know:

$$S \stackrel{*}{\Rightarrow} uAz$$

$$A \stackrel{*}{\Rightarrow} vAy$$

$$A \stackrel{*}{\Rightarrow} x$$

This string is also generated:

$$S \stackrel{*}{\Rightarrow} uAz \stackrel{*}{\Rightarrow} uvAyz \stackrel{*}{\Rightarrow} uvxyz$$

The original  $w = uv^1xy^1z$

We know:

$$S \xRightarrow{*} uAz$$

$$A \xRightarrow{*} vAy$$

$$A \xRightarrow{*} x$$

This string is also generated:

$$S \xRightarrow{*} uAz \xRightarrow{*} uvAyz \xRightarrow{*} uvvAyyz \xRightarrow{*} uvvxyyz$$

$$uv^2xy^2z$$

We know:

$$S \xRightarrow{*} uAz$$

$$A \xRightarrow{*} vAy$$

$$A \xRightarrow{*} x$$

This string is also generated:

$$\begin{aligned} S &\xRightarrow{*} uAz \xRightarrow{*} uvAyz \xRightarrow{*} uvvAyyz \xRightarrow{*} \\ &\xRightarrow{*} uvvvAyyyz \xRightarrow{*} uvvvxyyyz \end{aligned}$$

$$uv^3xy^3z$$

We know:

$$S \stackrel{*}{\Rightarrow} uAz$$

$$A \stackrel{*}{\Rightarrow} vAy$$

$$A \stackrel{*}{\Rightarrow} x$$

This string is also generated:

$$S \stackrel{*}{\Rightarrow} uAz \stackrel{*}{\Rightarrow} uvAyz \stackrel{*}{\Rightarrow} uvvAyyz \stackrel{*}{\Rightarrow}$$

$$\stackrel{*}{\Rightarrow} uvvvAyyyz \stackrel{*}{\Rightarrow} \dots$$

$$\stackrel{*}{\Rightarrow} uvvv \cdots vAy \cdots yyyz \stackrel{*}{\Rightarrow}$$

$$\stackrel{*}{\Rightarrow} uvvv \cdots vxy \cdots yyyz$$

$$uv^i xy^i z$$

Therefore, any string of the form

$$uv^i xy^i z \quad i \geq 0$$

is generated by the grammar  $G$




Therefore,


knowing that  $uvxyz \in L(G)$

we also know that  $uv^i xy^i z \in L(G)$

$$L(G) = L - \{\lambda\}$$




$$uv^i xy^i z \in L$$



Observation:  $|vxy| \leq m$

Since  $A$  is the last repeated variable



Observation:  $|vy| \geq 1$

Since there are no unit or  $\lambda$ -productions

# The Pumping Lemma:

For infinite context-free language  $L$

there exists an integer  $m$  such that

for any string  $w \in L$ ,  $|w| \geq m$

we can write  $w = uvxyz$

with lengths  $|vxy| \leq m$  and  $|vy| \geq 1$

and it must be:

$$uv^i xy^i z \in L, \quad \text{for all } i \geq 0$$

# Applications of The Pumping Lemma

# Non-context free languages

$$\{a^n b^n c^n : n \geq 0\}$$

Context-free languages

$$\{a^n b^n : n \geq 0\}$$

**Theorem:** The language

$$L = \{a^n b^n c^n : n \geq 0\}$$

is **not** context free

**Proof:** Use the Pumping Lemma  
for context-free languages

$$L = \{a^n b^n c^n : n \geq 0\}$$

Assume for contradiction that  $L$   
is context-free

Since  $L$  is context-free and infinite  
we can apply the pumping lemma



$$L = \{a^n b^n c^n : n \geq 0\}$$

Pumping Lemma gives a magic number  $m$   
such that:

Pick any string  $w \in L$  with length  $|w| \geq m$

We pick:  $w = a^m b^m c^m$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

We can write:  $w = uvxyz$

with lengths  $|vxy| \leq m$  and  $|vy| \geq 1$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Pumping Lemma says:

$$uv^i xy^i z \in L \quad \text{for all } i \geq 0$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$


We examine all the possible locations  
of string  $vxy$  in  $w$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 1:**  $vxy$  is within  $a^m$



$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 1:**  $v$  and  $y$  consist from only  $a$

$$\begin{array}{c}
 \begin{array}{ccc}
 m & m & m \\
 \underbrace{\hspace{1.5cm}} & \underbrace{\hspace{1.5cm}} & \underbrace{\hspace{1.5cm}} \\
 aaa\dots aaa & bbb\dots bbb & ccc\dots ccc \\
 \underbrace{\hspace{0.5cm}} & \underbrace{\hspace{0.5cm}} & \underbrace{\hspace{0.5cm}} \\
 u & vxy & z
 \end{array}
 \end{array}$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 1:** Repeating  $v$  and  $y$

$$k \geq 1$$

$$\begin{array}{c}
 \overbrace{a^{m+k} b^m c^m} \\
 \underbrace{a^{m+k} b^m c^m}_{\substack{u \quad v^2 xy^2 \quad z}}
 \end{array}$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 1:** From Pumping Lemma:  $uv^2xy^2z \in L$

$$k \geq 1$$

$$\begin{array}{c}
 \overbrace{a^{m+k}} \quad \overbrace{b^m} \quad \overbrace{c^m} \\
 aaaaaa\dots aaaaaa \quad bbb\dots bbb \quad ccc\dots ccc \\
 \underbrace{\quad} \quad \underbrace{\quad} \quad \underbrace{\quad} \\
 u \quad v^2xy^2 \quad z
 \end{array}$$



$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 1:** From Pumping Lemma:  $uv^2xy^2z \in L$

$$k \geq 1$$

However:  $uv^2xy^2z = a^{m+k} b^m c^m \notin L$


**Contradiction!!!**

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 2:**  $vxy$  is within  $b^m$



$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 2:** Similar analysis with case 1


$$\begin{array}{c}
 \begin{array}{ccc}
 m & m & m \\
 \underbrace{aaa \dots aaa} & \underbrace{bbb \dots bbb} & \underbrace{ccc \dots ccc} \\
 u & vxy & z
 \end{array}
 \end{array}$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 3:**  $vxy$  is within  $c^m$



$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 3:** Similar analysis with case 1


$$\begin{array}{c}
 m \qquad m \qquad m \\
 \underbrace{aaa\dots aaa} \quad \underbrace{bbb\dots bbb} \quad \underbrace{ccc\dots ccc} \\
 \underbrace{\hspace{10em}}_u \quad \underbrace{\hspace{10em}}_{vxy} \quad \underbrace{\hspace{10em}}_z
 \end{array}$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 4:**  $vxy$  overlaps  $a^m$  and  $b^m$




$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 4:** Possibility 1:  $v$  contains only  $a$   
 $y$  contains only  $b$



$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 4:** Possibility 1:  $v$  contains only  $a$

$k_1 + k_2 \geq 1$   $y$  contains only  $b$

$$\underbrace{aaa \dots a}_{m+k_1} \underbrace{bbb \dots b}_{m+k_2} \underbrace{ccc \dots c}_m$$

$$\underbrace{u}_{a} \quad \underbrace{v^2 xy^2}_{bb} \quad \underbrace{z}_{c}$$



$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 4:** From Pumping Lemma:  $uv^2xy^2z \in L$

$$k_1 + k_2 \geq 1$$

$$\underbrace{aaa \dots a}_{m+k_1} \underbrace{bbb \dots b}_{m+k_2} \underbrace{ccc \dots c}_m$$

$$\underbrace{u}_{u} \quad \underbrace{v^2 xy^2}_{v^2 xy^2} \quad \underbrace{z}_{z}$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 4:** From Pumping Lemma:  $uv^2xy^2z \in L$

$$k_1 + k_2 \geq 1$$

However:  $uv^2xy^2z = a^{m+k_1} b^{m+k_2} c^m \notin L$


**Contradiction!!!**

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 4:** Possibility 2:  $v$  contains  $a$  and  $b$   
 $y$  contains only  $b$




$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 4:** Possibility 2:  $v$  contains  $a$  and  $b$

$k_1 + k_2 + k \geq 1$        $y$  contains only  $b$



$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 4:** From Pumping Lemma:  $uv^2xy^2z \in L$

$$k_1 + k_2 + k \geq 1$$

$$\underbrace{aaa \dots aaa}_{m} \underbrace{abbaabb}_{k_1} \underbrace{bbbbb \dots bbb}_{m+k} \underbrace{ccc \dots ccc}_{m}$$

$$\underbrace{u}_{u} \underbrace{v^2 xy^2}_{v^2 xy^2} \underbrace{z}_{z}$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 4:** From Pumping Lemma:  $uv^2xy^2z \in L$

However:  $k_1 + k_2 + k \geq 1$

$$uv^2xy^2z = a^m b^{k_1} a^{k_2} b^{m+k} c^m \notin L$$


**Contradiction!!!**

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 4:** Possibility 3:  $v$  contains only  $a$   
 $y$  contains  $a$  and  $b$



$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 4:** Possibility 3:  $v$  contains only  $a$   
 $y$  contains  $a$  and  $b$

Similar analysis with Possibility 2




$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 5:**  $vxy$  overlaps  $b^m$  and  $c^m$



$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 5:** Similar analysis with case 4

$$\begin{array}{c}
 \begin{array}{ccc}
 m & m & m \\
 \underbrace{aaa \dots aaa} & \underbrace{bbb \dots bbb} & \underbrace{ccc \dots ccc} \\
 u & vxy & z
 \end{array}
 \end{array}$$

There are no other cases to consider

(since  $|vxy| \leq m$ , string  $vxy$  cannot

overlap  $a^m$ ,  $b^m$  and  $c^m$  at the same time)

In all cases we obtained a **contradiction**

**Therefore:** The original assumption that

$$L = \{a^n b^n c^n : n \geq 0\}$$

is context-free must be wrong

**Conclusion:**  $L$  is not context-free