More Applications of The Pumping Lemma

The Pumping Lemma:

For infinite context-free language L

there exists an integer m such that

for any string $w \in L, \quad |w| \geq m$

we can write $w = uvxyz$

with lengths $|vxy| \leq m$ and $|vy| \geq 1$

and it must be:

$$uv^i xy^i z \in L, \quad \text{for all } i \geq 0$$

Non-context free languages

$$\{a^n b^n c^n : n \geq 0\} \qquad \{vv : v \in \{a,b\}\}$$

Context-free languages

$$\{a^n b^n : n \geq 0\} \qquad \{ww^R : w \in \{a,b\}^*\}$$

Theorem: The language

$$L = \{vv : v \in \{a,b\}^*\}$$

is **not** context free

Proof: Use the Pumping Lemma for context-free languages

$$L = \{vv : v \in \{a,b\}^*\}$$

Assume for contradiction that L is context-free

Since L is context-free and infinite we can apply the pumping lemma

$$L = \{vv : v \in \{a,b\}^*\}$$

Pumping Lemma gives a magic number m such that:

Pick any string of L with length at least m

we pick: $a^m b^m a^m b^m \in L$

$$L = \{vv : v \in \{a,b\}^*\}$$

We can write: $a^m b^m a^m b^m = uvxyz$

with lengths $|vxy| \leq m$ and $|vy| \geq 1$

Pumping Lemma says:

$uv^i xy^i z \in L$ for all $i \geq 0$

$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \qquad |vxy| \leq m \qquad |vy| \geq 1$$

We examine **all** the possible locations of string vxy in $a^m b^m a^m b^m$

$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \qquad |vxy| \le m \qquad |vy| \ge 1$$

Case 1: vxy is within the first a^m

$$v = a^{k_1} \qquad y = a^{k_2} \qquad k_1 + k_2 \ge 1$$

$$\overbrace{a \ldots\ldots a}^{m}\ \overbrace{b \ldots\ldots b}^{m}\ \overbrace{a \ldots\ldots a}^{m}\ \overbrace{b \ldots\ldots b}^{m}$$

$$u \; v \; x \; y \qquad z$$

$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \qquad |vxy| \le m \qquad |vy| \ge 1$$

Case 1: vxy is within the first a^m

$$v = a^{k_1} \qquad y = a^{k_2} \qquad k_1 + k_2 \ge 1$$

$$\overbrace{a \ldots\ldots\ldots\ldots\ldots a}^{m+k_1+k_2}\;\overbrace{b \ldots\ldots b}^{m}\;\overbrace{a \ldots\ldots a}^{m}\;\overbrace{b \ldots\ldots b}^{m}$$

$$u \quad v^2 \quad x \quad y^2 \quad z$$

$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \qquad |vxy| \leq m \qquad |vy| \geq 1$$

$\boxed{\textbf{Case 1:} \quad vxy \text{ is within the first } a^m}$

$$a^{m+k_1+k_2} b^m a^m b^m = uv^2 xy^2 z \quad \notin L$$

$$k_1 + k_2 \geq 1$$

$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \qquad |vxy| \leq m \qquad |vy| \geq 1$$

$\boxed{\textbf{Case 1:} \quad vxy \text{ is within the first } a^m}$

$$a^{m+k_1+k_2} b^m a^m b^m = uv^2xy^2z \quad \notin L$$

However, from Pumping Lemma: $uv^2xy^2z \in L$

Contradiction!!!

$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \qquad |vxy| \le m \qquad |vy| \ge 1$$

Case 2: v is in the first a^m
y is in the first b^m

$$v = a^{k_1} \qquad y = b^{k_2} \qquad k_1 + k_2 \ge 1$$



$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \qquad |vxy| \leq m \qquad |vy| \geq 1$$

Case 2: v is in the first a^m

y is in the first b^m

$$v = a^{k_1} \qquad y = b^{k_2} \qquad k_1 + k_2 \geq 1$$

$$\overbrace{a\ldots\ldots\ldots\ldots a}^{m+k_1}\ \overbrace{b\ldots\ldots\ldots\ldots b}^{m+k_2}\ \overbrace{a\ldots\ldots a}^{m}\ \overbrace{b\ldots\ldots b}^{m}$$

$$u \quad v^2 \quad x \quad y^2 \quad z$$

$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \qquad |vxy| \leq m \qquad |vy| \geq 1$$

Case 2: v is in the first a^m

y is in the first b^m

$$a^{m+k_1} b^{m+k_2} a^m b^m = uv^2xy^2z \quad \notin L$$

$$k_1 + k_2 \geq 1$$

$$L=\{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \qquad |vxy| \le m \qquad |vy| \ge 1$$

Case 2: v is in the first a^m
y is in the first b^m

$$a^{m+k_1} b^{m+k_2} a^m b^m = uv^2xy^2z \quad \notin L$$

However, from Pumping Lemma: $uv^2xy^2z \in L$

Contradiction!!!

$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \qquad |vxy| \le m \qquad |vy| \ge 1$$

Case 3: v overlaps the first $a^m b^m$
y is in the first b^m

$$v = a^{k_1} b^{k_2} \qquad y = b^{k_3} \qquad k_1, k_2 \ge 1$$



$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \qquad |vxy| \le m \qquad |vy| \ge 1$$

Case 3: v overlaps the first $a^m b^m$
y is in the first b^m

$$v = a^{k_1} b^{k_2} \qquad y = b^{k_3} \qquad k_1, k_2 \ge 1$$

$$\overbrace{a \ldots\ldots a}^{m}\ \overbrace{b \ldots b}^{k_2}\ \overbrace{a \ldots a}^{k_1}\ \overbrace{b \ldots\ldots\ldots b}^{m+k_3}\ \overbrace{a \ldots\ldots a}^{m}\ \overbrace{b \ldots\ldots b}^{m}$$

$$u \qquad v^2 \qquad x \qquad y^2 \qquad z$$

$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \qquad |vxy| \leq m \qquad |vy| \geq 1$$

Case 3: v overlaps the first $a^m b^m$

y is in the first b^m

$$a^m b^{k_2} a^{k_1} b^{m+k_3} a^m b^m = uv^2 xy^2 z \quad \notin L$$

$$k_1, k_2 \geq 1$$

$$L=\{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \qquad |vxy| \leq m \qquad |vy| \geq 1$$

Case 3: v overlaps the first $a^m b^m$
y is in the first b^m

$$a^m b^{k_2} a^{k_1} b^{k_3} a^m b^m = uv^2xy^2z \quad \notin L$$

However, from Pumping Lemma: $uv^2xy^2z \in L$

Contradiction!!!

$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \qquad |vxy| \le m \qquad |vy| \ge 1$$

Case 4: v in the first a^m

y Overlaps the first $a^m b^m$

Analysis is similar to case 3



Other cases: vxy is within $a^m \boxed{b^m} a^m b^m$

or

$a^m b^m \boxed{a^m} b^m$

or

$a^m b^m a^m \boxed{b^m}$

Analysis is similar to case 1:

$$\boxed{a^m} b^m a^m b^m$$

More cases: vxy overlaps $a^m \boxed{b^m a^m} b^m$

or

$a^m b^m \boxed{a^m b^m}$

Analysis is similar to cases 2,3,4:

$\boxed{a^m b^m} a^m b^m$

There are no other cases to consider

Since $|vxy| \leq m$, it is impossible vxy to overlap:

$$\boxed{a^m b^m a^m} b^m$$

nor

$$a^m \boxed{b^m a^m b^m}$$

nor

$$\boxed{a^m b^m a^m b^m}$$

In all cases we obtained a contradiction

Therefore: The original assumption that

$$L = \{vv : v \in \{a,b\}^*\}$$

is context-free must be wrong

Conclusion: L is not context-free

Non-context free languages

$$\{a^n b^n c^n : n \geq 0\}$$

$$\{ww : w \in \{a, b\}\}$$

$$\{a^{n!} : n \geq 0\}$$

Context-free languages

$$\{a^n b^n : n \geq 0\}$$

$$\{ww^R : w \in \{a, b\}^*\}$$

Theorem: The language

$$L = \{a^{n!} : n \geq 0\}$$

is **not** context free

Proof: Use the Pumping Lemma for context-free languages

$$L = \{a^{n!} : n \geq 0\}$$

Assume for contradiction that L is context-free

Since L is context-free and infinite we can apply the pumping lemma

$$L = \{a^{n!} : n \geq 0\}$$

Pumping Lemma gives a magic number m such that:

Pick any string of L with length at least m

we pick: $a^{m!} \in L$

$$L = \{a^{n!} : n \geq 0\}$$

We can write: $a^{m!} = uvxyz$

with lengths $|vxy| \leq m$ and $|vy| \geq 1$

Pumping Lemma says:

$uv^i xy^i z \in L$ for all $i \geq 0$

$$L = \{a^{n!} : n \geq 0\}$$

$$a^{m!} = uvxyz \qquad |vxy| \leq m \qquad |vy| \geq 1$$

We examine **all** the possible locations of string vxy in $a^{m!}$

There is only one case to consider

$$L = \{a^{n!} : n \geq 0\}$$

$$a^{m!} = uvxyz \qquad |vxy| \leq m \qquad |vy| \geq 1$$



$$v = a^{k_1} \qquad y = a^{k_2} \qquad 1 \leq k_1 + k_2 \leq m$$

$$L = \{a^{n!} : n \geq 0\}$$

$$a^{m!} = uvxyz \qquad |vxy| \leq m \qquad |vy| \geq 1$$



$$v = a^{k_1} \qquad y = a^{k_2} \qquad 1 \leq k_1 + k_2 \leq m$$

$$L = \{a^{n!} : n \geq 0\}$$

$$a^{m!} = uvxyz \qquad |vxy| \leq m \qquad |vy| \geq 1$$

$$\overbrace{a\ \ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ \ a}^{m!+k} \qquad k = k_1 + k_2$$

$$u \quad v^2 \quad x \quad y^2 \quad z$$

$$v = a^{k_1} \qquad y = a^{k_2} \qquad 1 \leq k \leq m$$

$$L = \{a^{n!} : n \geq 0\}$$

$$a^{m!} = uvxyz \qquad |vxy| \leq m \qquad |vy| \geq 1$$

$$a^{m!+k} = uv^2xy^2z$$

$$1 \leq k \leq m$$

Since $1 \leq k \leq m$, for $m \geq 2$ we have:

$$m! + k \leq m! + m$$

$$< m! + m!m$$

$$= m!(1 + m)$$

$$= (m + 1)!$$

$$\Downarrow$$

$$m! < m! + k < (m + 1)!$$

$$L = \{a^{n!} : n \geq 0\}$$

$$a^{m!} = uvxyz \qquad |vxy| \leq m \qquad |vy| \geq 1$$

$$m! < m! + k < (m+1)!$$

$$\Downarrow$$

$$a^{m!+k} = uv^2xy^2z \quad \notin L$$

$$L = \{a^{n!} : n \geq 0\}$$

$$a^{m!} = uvxyz \qquad |vxy| \leq m \qquad |vy| \geq 1$$

However, from Pumping Lemma: $uv^2xy^2z \in L$

$$a^{m!+k} = uv^2xy^2z \quad \notin L$$

Contradiction!!!

We obtained a contradiction

Therefore: The original assumption that

$$L = \{a^{n!} : n \geq 0\}$$

is context-free must be wrong

Conclusion: L is not context-free

Non-context free languages

$$\{a^n b^n c^n : n \geq 0\}$$

$$\{ww : w \in \{a,b\}\}$$

$$\{a^{n^2} b^n : n \geq 0\}$$

$$\{a^{n!} : n \geq 0\}$$

Context-free languages

$$\{a^n b^n : n \geq 0\}$$

$$\{ww^R : w \in \{a,b\}^*\}$$

Theorem: The language

$$L = \{a^{n^2} b^n : n \geq 0\}$$

is **not** context free

Proof: Use the Pumping Lemma for context-free languages

$$L = \{a^{n^2} b^n : n \geq 0\}$$

Assume for contradiction that L is context-free

Since L is context-free and infinite we can apply the pumping lemma

$$L = \{a^{n^2} b^n : n \geq 0\}$$

Pumping Lemma gives a magic number m such that:

Pick any string of L with length at least m

we pick: $a^{m^2} b^m \in L$

$$L = \{a^{n^2} b^n : n \geq 0\}$$

We can write: $a^{m^2} b^m = uvxyz$

with lengths $|vxy| \leq m$ and $|vy| \geq 1$

Pumping Lemma says:

$uv^i xy^i z \in L$ for all $i \geq 0$

$$L = \{a^{n^2} b^n : n \geq 0\}$$

$$a^{m^2} b^m = uvxyz \qquad |vxy| \leq m \qquad |vy| \geq 1$$

We examine **<u>all</u>** the possible locations

of string vxy in $a^{m^2} b^m$

$$L = \{a^{n^2} b^n : n \geq 0\}$$

$$a^{m^2} b^m = uvxyz \qquad |vxy| \leq m \qquad |vy| \geq 1$$

Most complicated case: v is in a^m

y is in b^m



$$L = \{a^{n^2} b^n : n \geq 0\}$$

$$a^{m^2} b^m = uvxyz \qquad |vxy| \leq m \qquad |vy| \geq 1$$

$$v = a^{k_1} \qquad y = b^{k_2} \qquad 1 \leq k_1 + k_2 \leq m$$



$$L = \{a^{n^2} b^n : n \geq 0\}$$

$$a^{m^2} b^m = uvxyz \qquad |vxy| \leq m \qquad |vy| \geq 1$$

Most complicated sub-case: $k_1 \neq 0$ and $k_2 \neq 0$

$$v = a^{k_1} \qquad y = b^{k_2} \qquad 1 \leq k_1 + k_2 \leq m$$



$$L = \{a^{n^2} b^n : n \geq 0\}$$

$$a^{m^2} b^m = uvxyz \qquad |vxy| \leq m \qquad |vy| \geq 1$$

Most complicated sub-case: $k_1 \neq 0$ and $k_2 \neq 0$

$$v = a^{k_1} \qquad y = b^{k_2} \qquad 1 \leq k_1 + k_2 \leq m$$

$$\overbrace{a \ldots\ldots\ldots\ldots\ldots a}^{m^2 - k_1}\ \overbrace{b \ldots b}^{m - k_2}$$

$$u \qquad v^0 \qquad x \qquad y^0 \qquad z$$

$$L = \{a^{n^2} b^n : n \geq 0\}$$

$$a^{m^2} b^m = uvxyz \qquad |vxy| \leq m \qquad |vy| \geq 1$$

Most complicated sub-case: $k_1 \neq 0$ and $k_2 \neq 0$

$$v = a^{k_1} \qquad y = b^{k_2} \qquad 1 \leq k_1 + k_2 \leq m$$

$$a^{m^2 - k_1} b^{m - k_2} = uv^0 xy^0 z$$

$k_1 \neq 0$ and $k_2 \neq 0$ $\quad 1 \leq k_1 + k_2 \leq m$

$$\Downarrow$$

$$(m - k_2)^2 \leq (m-1)^2$$

$$= m^2 - 2m + 1$$

$$< m^2 - k_1$$

$$\Downarrow$$

$$m^2 - k_1 \neq (m - k_2)^2$$

$$L = \{a^{n^2} b^n : n \geq 0\}$$

$$a^{m^2} b^m = uvxyz \qquad |vxy| \leq m \qquad |vy| \geq 1$$

$$m^2 - k_1 \neq (m - k_2)^2$$

$$\Downarrow$$

$$a^{m^2 - k_1} b^{m - k_2} = uv^0 xy^0 z \quad \notin L$$

$$L = \{a^{n^2} b^n : n \geq 0\}$$

$$a^{m^2} b^m = uvxyz \qquad |vxy| \leq m \qquad |vy| \geq 1$$

However, from Pumping Lemma: $uv^0xy^0z \in L$

$$a^{m^2-k_1} b^{m-k_2} = uv^0xy^0z \notin L$$

Contradiction!!!

When we examine the rest of the cases we also obtain a contradiction

In all cases we obtained a contradiction

Therefore: The original assumption that

$$L = \{a^{n^2} b^n : n \geq 0\}$$

is context-free must be wrong

Conclusion: L is not context-free